

Final[®] RODENTICIDE



New Rodenticide with Brodifacoum

Bell Laboratories introduces its newest single-feeding, anticoagulant rodenticide, FINAL Rodenticide, with the powerful active ingredient Brodifacoum. FINAL combines the superior formulating expertise of Bell Laboratories with the strongest anticoagulant available on the market today. Containing more than 16 human food-grade inert ingredients, FINAL is more palatable than other Brodifacoum baits on the market.

Effective on rats and mice, FINAL with Brodifacoum is especially successful in cleaning out tough mouse infestations where populations are difficult to control or where resistance is suspected.

Packaging: 10 lb. bulk plastic pail – EPA REG. NO. 12455-90
25 lb. bulk plastic pail – EPA REG. NO. 12455-90
100 x 25 g. place pacs – EPA REG. NO. 12455-91
200 x 25 g. place pacs – EPA REG. NO. 12455-91

Specimen Label

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice in and around homes, industrial, commercial, agricultural and public buildings, and similar man-made structures. FINAL Rodenticide may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. This product may also be used in alleys. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait. When used in USDA inspected plants, this product must be applied in tamper-resistant bait stations.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

(Continued at the top of the right panel)

Final[®] RODENTICIDE

KILLS RATS AND MICE

Kills Warfarin Resistant Norway Rats and House Mice

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENTS: Brodifacoum;
3-[3-(4'-Bromo-[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one..... 0.005%
INERT INGREDIENTS* 99.995%
TOTAL 100.000%
*Contains Denatonium Benzoate

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

If on Skin: Wash with plenty of soap and water. Get medical attention if symptoms persist.

If in Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K, intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

(See right side panel for additional precautionary statements)

For 24-hour emergency medical advice, call 1-800-228-5635 ext. 311.

Manufactured by:



Bell Laboratories, Inc.
Madison, WI 53704 U.S.A.

EPA EST. NO. 12455-WI-1 EPA REG. NO. 12455-90

DIRECTIONS FOR USE (Continued from left panel.)

APPLICATION DIRECTIONS:

RATS: Apply 3 to 16 ounces of bait (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

HOUSE MICE: Apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait per placement at 8- to 12- foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

RATS/MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling bait.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store only in original container in a dry area inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty pail. Then dispose of empty pail in sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.